



TITLE:

# Modulation of Ig $\beta$ is essential for the B cell selection in germinal center

AUTHOR(S):

Todo, Kagefumi; Koga, Orie; Nishikawa, Miwako; Hikida, Masaki

---

CITATION:

Todo, Kagefumi ...[et al]. Modulation of Ig $\beta$  is essential for the B cell selection in germinal center. Scientific Reports 2015, 5: 10303.

ISSUE DATE:

2015-05-18

URL:

<http://hdl.handle.net/2433/212467>

RIGHT:

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

# SCIENTIFIC REPORTS

OPEN

## Modulation of Ig $\beta$ is essential for the B cell selection in germinal center

Kagefumi Todo, Orié Koga, Miwako Nishikawa & Masaki Hikida

Received: 20 October 2014

Accepted: 08 April 2015

Published: 18 May 2015

The positive and negative selection of antigen-reactive B cells take place in the germinal center (GC) during an immune responses. However, the precise molecular mechanisms underlying these selection machineries, including the involvement of antigen receptor signaling molecules, remain to be elucidated. We found that expression levels of Ig $\alpha$  and Ig $\beta$ , which are the essential components of B cell antigen-receptor complex, were differentially regulated in GC B cells and that the expression of Ig $\beta$  was more prominently down-regulated in a portion of GC B cells. The suppression of Ig $\beta$  down-regulation reduced the number of GL7<sup>+</sup> GC B cells and the affinity maturation in T-dependent responses was markedly impaired. In addition, the disease phenotypes in autoimmune-prone mice were ameliorated by blocking of Ig $\beta$  down-regulation. These results suggest that Ig $\beta$  down-regulation is involved in the normal positive selection in GC and the accumulation of autoreactive B cells in autoimmune-prone mice.

The B cell antigen receptor (BCR) is a protein complex that consists of a membrane-bound immunoglobulin (Ig) molecule and the signal transducer, an Ig $\alpha$ /Ig $\beta$  hetero-dimer, and other signaling molecules<sup>1–3</sup>. It is well known that the Ig $\alpha$ /Ig $\beta$  hetero-dimer is required for the expression of membrane-bound Ig $\mu$  chains on the surface of preB cells<sup>1,4,5</sup>. Furthermore, Ig $\mu$  chains expressed on the cell surface of B lineage cells in association with Ig $\alpha$ /Ig $\beta$  hetero-dimer play essential roles in both differentiation and survival of preB cells and mature B cells<sup>6,7</sup>. In addition, it has been shown that cell surface expression of the Ig $\alpha$ /Ig $\beta$  hetero-dimer not only supports the expression of cell surface Ig $\mu$  chains, but also the signal through this complex is further required for the differentiation and survival of B lineage cells<sup>8–10</sup>. Hence, it has been widely believed that both Ig $\alpha$  and Ig $\beta$  are expressed in B lineage cells during all maturation stages.

After completion of differentiation, mature B cells participate in the humoral immune responses. One of the hallmarks of the humoral immune response is the formation of germinal centers (GCs) following the activation of B cells by an antigen under the influence of T cells<sup>11–13</sup>. It is widely known that GC B cells can be classified into two compartments namely centroblasts and centrocytes. Centroblasts are observed in the dark zone and they lack or express only low levels of surface Ig. These cells proceed with somatic hypermutation of their antibody variable genes and proliferate rapidly, which contribute to the clonal expansion. In contrast, centrocytes are relatively small non-dividing cells with surface Ig in the light zone where positive and negative selection take place<sup>14</sup>. A combination of somatic hypermutation, clonal expansion, and selection leads a part of GC B cells to acquire a BCR with higher affinity for the antigen, which results in the affinity maturation of serum antibodies. It has been widely shown that a portion of GC B cells, mainly consisting of centroblasts, reduces their surface BCR expression during these processes. Thus, it can easily be predicted that BCR-associating molecules, including Ig $\alpha$  and Ig $\beta$ , are down-regulated in these cells. Indeed, it has been reported that expression of both Ig $\alpha$  and Ig $\beta$  was down-regulated in the germinal center (GC) B cells<sup>15–17</sup>. However, it has not been determined whether the modulation of these signaling molecules has as-yet-unknown physiological roles or simply reflects BCR down-regulation.

Center for Innovation in Immunoregulative Technology and Therapeutics, Graduate School of Medicine, Kyoto University Yoshidakonocho, Sakyo-ku, Kyoto 606-8501, Japan. Correspondence and requests for materials should be addressed to M.H. (email: hikida@ak.med.kyoto-u.ac.jp)

In this study, we demonstrated that expression levels of Ig $\alpha$  and Ig $\beta$ , were differentially regulated in GC B cells and that the expression of Ig $\beta$  was more prominently down-regulated in a part of GC B cells. Furthermore, this down-regulation of Ig $\beta$  is involved both in the effective positive selection in GC B cells and the accumulation of autoreactive B cells in autoimmune-prone mice.


## Results

**The expression of Ig $\beta$  is down-regulated in GC B cells.** It has been reported that Ig $\beta$  is ubiquitously expressed in both immature and mature B cells. However, it has not been fully investigated whether Ig $\beta$  is also expressed constantly in B cells during immune responses, such as in GC B cells. To clarify this point, we initially analyzed the expression of Ig $\beta$  in the spleen from immunized mice by immunohistochemical staining. Ten days post immunization with NP-CGG, PNA<sup>+</sup>CD38<sup>−</sup> GCs were clearly detected (Fig. 1a). When compared with the follicular B cells, B cells in GCs were only weakly stained by anti-Ig $\beta$  antibodies (Fig. 1a). Spleen cells from immunized mice were further analyzed by flow cytometer to confirm the down-regulation of cell surface Ig $\beta$ . As shown in Fig. 1b, naïve B cells (B220<sup>+</sup>CD38<sup>+</sup>IgM<sup>+</sup>) expressed Ig $\beta$  at high levels (MFI =  $9.7 \times 10^3$ ), as expected. In contrast, the levels of Ig $\beta$  decreased in GC B cells identified with either surface markers CD38 or GL7 (B220<sup>+</sup>CD38<sup>−</sup>: MFI =  $2.1 \times 10^3$  and B220<sup>+</sup>GL7<sup>+</sup>: MFI =  $3.5 \times 10^3$ ). Quantitative RT-PCR revealed that the level of Ig $\beta$  mRNA in non-apoptotic GC B cells was reduced by approximately 50% of that in naïve B cells (Fig. 1c), suggesting that the cell surface expression of Ig $\beta$  is controlled at the transcription level to some extent. According to a previous report<sup>15</sup> and the present data (see below), the expression of Ig $\beta$  in GC B cells clearly correlated with the down-modulation of IgM. Thus, to clarify whether this down-regulation of Ig $\beta$  in GC B cells simply reflects the decrease of surface BCR expression, we examined the levels of Ig $\beta$  mRNA following the endocytosis of BCR *in vitro* (Fig. 1d). In comparison with the decreased surface IgM levels after BCR cross-linking, expression of Ig $\beta$  mRNA was sustained at a constant level. Although cell surface level of Ig $\beta$  correlates well with the endocytosis of IgM, these results suggest that there is an additional regulatory mechanism for the down-regulation of Ig $\beta$  mRNA *in vivo* independent from BCR endocytosis induced by simple cross-linking (see below).

**GC B cells can be classified into several stages based on their surface Ig $\beta$  expression.** When B220<sup>+</sup>CD38<sup>−</sup> B cells from immunized mice were analyzed for GL7 expression, the resulting population could be classified into GL7<sup>+</sup> and GL7<sup>−</sup> subsets (Fig. 2a, left). Both of these subsets were IgD<sup>−</sup>, another GC B cell marker, suggesting that these subsets are both GC B cells (Fig. 2b). The B220<sup>+</sup>CD38<sup>−</sup>GL7<sup>+</sup> B cells formed a unique subset whose expression level of Ig $\beta$  was 1/4 to 1/5 of that in B220<sup>+</sup>CD38<sup>+</sup> naïve B cells (Fig. 2a, right bottom). In contrast, CD38<sup>−</sup>GL7<sup>−</sup> B cells could be further divided into three subpopulations based on their surface level of Ig $\beta$ . Approximately 23% of the CD38<sup>−</sup>GL7<sup>−</sup> cells expressed Ig $\beta$  at a level similar to that in naïve B cells (Ig $\beta$ -Hi cells), 43% of CD38<sup>−</sup>GL7<sup>−</sup> B cells expressed Ig $\beta$  at a level of 1/8 to 1/10 of the level in naïve B cells (Ig $\beta$ -Int cells), and 34% of these cells did not express or, at most, expressed low levels of Ig $\beta$  (Ig $\beta$ -Lo cells) (Fig. 2a, right bottom). In contrast, the cell surface expression of Ig $\alpha$  was only slightly down-regulated in the CD38<sup>−</sup>GL7<sup>−</sup> B cell subset (MFI = 42.2) as compared with naïve B cells (MFI = 55.7), and the level of Ig $\alpha$  was not down-regulated in CD38<sup>−</sup>GL7<sup>+</sup> B cells (MFI = 65.8) (Fig. 2c). These results are consistent with the previous histological analysis of human lymph nodes reported by others<sup>17</sup>, strongly suggesting that expression of Ig $\alpha$  and Ig $\beta$  are differentially regulated in GC B cells.

To further confirm that this B220<sup>+</sup>CD38<sup>−</sup>GL7<sup>−</sup> population represents the subsets of GC B cells, we analyzed the expression of Bcl6 and AID (Fig. 2d, left). The expression of Ig $\beta$  mRNA correlated well with the cell surface Ig $\beta$  level, which again suggests that Ig $\beta$  expression in GC B cells is regulated primarily at the transcription level. Bcl6 and AID were highly expressed in the Ig $\beta$ -Hi, Ig $\beta$ -Int and GL7<sup>+</sup> cells. These results clearly demonstrate that the Ig $\beta$ -Hi and Ig $\beta$ -Int B cell subsets were indeed GC B cells. We also examined the expression level of Fas, which has been reported to be expressed in activated and GC B cells<sup>18,19</sup> (Fig. 2d, right). High levels of Fas expression were detected in a majority of the GL7<sup>+</sup> B cells. Among the GL7<sup>−</sup> subpopulation, the Ig $\beta$ -Int population significantly up-regulated Fas expression suggesting that they are either a part of the GC B cells or at least of the activated B cells. We also checked the expression level of PD-1 in these cells and found that these Ig $\beta$ -Int cells expressed only quite a low level of PD-1 (Fig. S1a), suggesting that Ig $\beta$ -Int cells are a distinct population from the reported memory B cell fraction<sup>20</sup>. In addition, this B220<sup>+</sup>CD38<sup>−</sup>GL7<sup>−</sup> population was confirmed to be CD138<sup>−</sup> indicating that they were not plasma cell committed cells (data not shown). We further examined the expression levels of several known markers for GC B cells and memory B cells in B220<sup>+</sup>CD38<sup>−</sup>GL7<sup>−</sup> B cells and found that Ig $\beta$ -Hi cells do not express memory surface markers (Fig. S1b).


To identify the proliferating cells among the aforementioned GC subpopulations, spleen cells from immunized mice were stained with the membrane-permeable DNA staining dye Hoechst33342 (Fig. 2e). Most IgM<sup>+</sup>CD38<sup>+</sup>B cells (>90%) were in the G0/G1 phase and a small numbers of cell were in the S and G2/M phase (Fig. 2e, top). In contrast, a significant number of the GL7<sup>+</sup>B cells (6%-7%) were in the S + G2/M phase (Fig. 2e, bottom right). On the other hand, most of the cells in the S + G2/M phase were Ig $\beta$ -Int cells belonging to the B220<sup>+</sup>CD38<sup>−</sup>GL7<sup>−</sup> population. Considering that GC B cells consisted of similar number of GL7<sup>+</sup> and GL7<sup>−</sup> populations (Fig. 2a, right bottom), these results may suggest that the major proliferating cells in the GC belong to the GL7<sup>−</sup>Ig $\beta$ -Int B cell population (Fig. 2e, bottom left).



**Figure 1.** Down-regulated Igβ expression in the germinal center. **(a)** Frozen sections of the spleen from B6 mice, 10 days after the immunization with NP-CGG precipitated in alum, were stained with PNA (green) and antibodies to Igβ (red) and CD38 (blue). **(b)** Naïve B cells (shaded), B220<sup>+</sup>CD38<sup>-</sup> cells (blue line) and B220<sup>+</sup>GL7<sup>+</sup> cells (red line) were analyzed for the expression level of Igβ. **(c)** Quantitative RT-PCR of Igβ mRNA in sorted naïve B cells (B220<sup>+</sup>IgM<sup>+</sup>CD38<sup>+</sup>) and GC B cells (B220<sup>+</sup>CD38<sup>-</sup>) from B6 mice 10 days after the immunization with NP-CGG. Cells were stained with propidium iodide to exclude dead cells and apoptotic cells. Data were normalized to the expression levels of the β-actin transcript. **(d)** Surface IgM in mouse splenic B cells was cross-linked with 10 μg/ml anti-IgM F(ab')<sub>2</sub> to induce endocytosis for indicated periods of time and remaining levels of cell surface IgM were analyzed (**left**). Expression levels of Igβ mRNA were analyzed by RT-qPCR in B cells whose surface IgM was down-modulated by cross-linking for the indicated periods (**right**).


Furthermore, it was shown that most of the Igβ<sup>Lo</sup> cells belonged to the sub-G0 phase indicating that these cells are apoptotic (Fig. 2e, bottom left).

We next examined the expression of surface Igβ in combination with surface IgG1 instead of GL7. The level of Igβ changed dramatically along with the acquisition of surface IgG1 in the B220<sup>+</sup>CD38<sup>-</sup> B cells, which prompted the need to analyze class switching in each population in detail (Fig. 3a, bottom).



**Figure 2.** Classification of GC B cells by expression level of Igβ. **(a)** Flow cytometric analysis of splenocytes from immunized B6 mice. B220+B cells (left) cells were further gated to B220+CD38+ cells (**top right**) and B220+CD38- cells (**bottom right**) and analyzed for the expression levels of Igβ and GL7. The numbers indicate the percentage of cells in each square, and the number in the parenthesis indicate the percentage of cells in each square within the CD38-GL7- cells. **(b)** Suppressed IgD expression in B220+CD38-GL7- and B220+CD38-GL7+B cells. **(c)** Expression of cell surface Igα in GC B cells. Cell surface expression levels of Igα and GL7 were examined in B220+CD38+naive B cells (**top**) and B220+CD38- GC B cells (**bottom**) by staining with anti-GL7 mAb in combination with rabbit polyclonal antibody raised against extracellular portion of Igα. Numbers indicate the percentages of the cells within the gates. **(d)** Transcription levels of Igβ, Bcl6 and AID in GC subsets (**left**) and surface Fas expression in B220+CD38-GL7- and B220+CD38-GL7+GC subsets (**right**). RT-qPCR were carried out in sorted B220+CD38-GL7-IgβHi cells (Igβ-Hi), B220+CD38-GL7-IgβInt cells (Igβ-Int), B220+CD38-GL7-IgβLo cells (Igβ-Lo) and B220+CD38-GL7+ cells (GL7+) 10 days after the immunization. **(e)** Cell cycle analysis of GC B cells. Splenocytes from immunized B6 mice were stained with Hoechst 33342 dye in combination with antibodies to Igβ and indicated surface markers, followed by FACS analysis. Naive B cells (**top**), B220+CD38-GL7- (**bottom left**) and B220+CD38-GL7+B cells (**bottom right**) were analyzed.





**Figure 3.** Class switching events in GC B cells (a) Expression of IgG1 and Ig $\beta$  in B220<sup>+</sup>IgM<sup>+</sup>CD38<sup>+</sup> naive B cells (top) and B220<sup>+</sup>CD38<sup>-</sup> GC B cells (bottom). (b) Expression levels of germline transcript and surface expression of IgM and IgG1 in the GC B cell subsets. Expression of IgM germline transcript (GLT- $\mu$ ), IgG1 germline transcript (GLT- $\gamma$ 1) and IgG1 postswitch transcript were assessed by RT-qPCR in each population, respectively (top). Cell surface expression of IgM and IgG1 in each GC subset was examined (bottom). (c) Frequency of somatic hypermutation in VH186.2 gene of GC B cell subsets. Number of nucleic acid mutations (left) and amino acid mutations (right) are shown.

Thus, the expression of immunoglobulin mRNA was tested to clarify the class switching status of both GL7<sup>−</sup> and GL7<sup>+</sup> subsets identified in Fig. 2a. As the result, high levels of I $\mu$  germline transcripts were detected in the GL7<sup>−</sup> Ig $\beta$ -Hi and Ig $\beta$ -Int cells. Since it has been reported that I $\mu$  germline transcripts were constitutively transcribed from the IgM locus in the germline configuration<sup>21</sup> (Fig. 3b, top), it is apparent that these populations include IgM-bearing non class-switched B cells as a major population. Comparable levels of I $\gamma$ 1 germline transcripts and  $\gamma$ 1 post-switch transcripts were detected in Ig $\beta$ -Int cells when compared with GL7<sup>+</sup> cells (Fig. 3b, top). Since it has been reported that expression of germline transcripts take place prior to DNA rearrangement<sup>22</sup>, these results suggest that class switching from IgM to IgG1 may take place in both the GL7<sup>−</sup> Ig $\beta$ -Int population and the GL7<sup>+</sup> stage. In accordance with these data, nearly 90% of the cells in the GL7<sup>−</sup> Ig $\beta$ -Hi population expressed IgM on their surface, including the population beginning to down-regulate IgM (Fig. 3b, bottom left). In contrast, around 60% of the Ig $\beta$ -Int population expressed only a low level of IgM and a portion of these cells (8.7%) begin to express a low, although significant, level of IgG1 on their surface and IgG1 single-positive cells were clearly detected in the GL7<sup>+</sup> population (Fig. 3b, bottom right). According to a previous report<sup>23</sup>, the stimulation of IgM<sup>+</sup>B cells with anti-CD40 or CD40 ligand together with IL-4 induced only GL7<sup>+</sup>B cells concomitantly with class switching to IgG1. A straightforward interpretation for the previous and the aforementioned results is that, in addition to the population that became IgM<sup>−</sup>GL7<sup>+</sup> class-switched cells following the IgM<sup>+</sup>GL7<sup>+</sup> population, a significant portion of IgM<sup>+</sup>GL7<sup>−</sup> GC B cells might differentiate to IgM<sup>−</sup>GL7<sup>+</sup> class-switched cells via the IgM<sup>dull</sup>GL7<sup>−</sup> Ig $\beta$ -Int stage. Indeed, both IgM<sup>+</sup>GL7<sup>+</sup> and IgM<sup>dull</sup>GL7<sup>−</sup> GC B cell subsets were detected (Fig. S2). This result may suggest the existence of a novel class-switching pathway distinct from *in vitro* class-switching where B cells differentiate from IgM<sup>+</sup>GL7<sup>−</sup> to IgM<sup>dull</sup>GL7<sup>+</sup> switched B cells through IgM<sup>+</sup>GL7<sup>+</sup>B cells. In the *in vitro* system, it was shown that the considerable number of somatic hypermutations in the Ig gene were detected only when AID was over expressed<sup>23</sup>. This is consistent with our results that Ig $\beta$ -Int cells expressed higher level of AID (Fig. 2d).

To approach the speculation that Ig $\beta$ -Int GL7<sup>−</sup> B cells may contribute to positive selection in the GC, the frequency of somatic hypermutation in each subset was analyzed (Fig. 3c). On day 10 post-immunization of NP-CGG, nucleotide mutations in VH186.2 were readily detected in Ig $\beta$ -Hi cells and the frequency slightly increased in the Ig $\beta$ -Int and GL7<sup>+</sup> cells, although the increase was not statistically significant (Fig. 3c, left). In parallel, amino acid replacements were significantly accumulated in the GL7<sup>+</sup> cells as compared with Ig $\beta$ -Hi cells. These results suggest that somatic hypermutations accumulate throughout the CD38<sup>−</sup> stage, including both the GL7<sup>−</sup> and GL7<sup>+</sup> populations, and that affinity maturation may proceed in the Ig $\beta$ -Int cells (Fig. 3c, right).


**Blocking of Ig $\beta$  down-regulation in GC B cells impairs GC formation, efficient Ig production and affinity maturation.** To further evaluate the physiological roles of Ig $\beta$  down-regulation, B cell-specific Ig $\beta$ -transgenic (Ig $\beta$ -Tg) mice were generated (Fig. S3). Between the established two Tg lines, the line with higher expression of exogenous Ig $\beta$  (line #1) was primarily analyzed. Over all differentiation of B cells in these mice was normal, except an increase in B220<sup>dull</sup>CD43<sup>−</sup> pre-B cells was observed (Fig. S3d). Furthermore, there were no significant differences in spontaneous Ig titers between transgenic mice and wild-type littermate controls (Fig. S3e). In addition, endocytosis of IgM complex by cross-linking of BCR was not affected in these transgenic mice (data not shown).

Frequency of Ig $\beta$  down-regulated cells in the B220<sup>+</sup>CD38<sup>−</sup>GL7<sup>−</sup> B cells from the immunized Ig $\beta$ -Tg mice (28.3%) decreased significantly compared to that in the wild-type littermate controls (74.9%), which is consistent with the results obtained in C57BL/6 mice (Figs. 4a and 2a). This finding indicates that the transgene was expressed in the expected manner, allowing us to analyze the function of Ig $\beta$  down-regulation in GC B cells.

Next, the frequency of the GC B cell subsets defined above was enumerated. Normal GC formation was observed by histochemical analysis (Fig. S3f). The frequency of B220<sup>+</sup>CD38<sup>−</sup>GL7<sup>+</sup> cells in the immunized spleen of the Ig $\beta$ -Tg mice decreased to approximately 56% of that of wild-type mice (0.69% to 0.24%) (Fig. 4b top). In addition, the cell cycle status of the B220<sup>+</sup>CD38<sup>−</sup>GL7<sup>−</sup> B cells was examined and the proportion of proliferating cells was reduced from 10.8% to 4.4% (Fig. 4b, bottom).

These results suggest the involvement of Ig $\beta$  down-regulation in the differentiation of GC B cells to proliferating cells, which may affect clonal expansion and subsequent affinity maturation. To investigate this possibility, NP-specific Ig titers and their relative affinity in Ig $\beta$ -Tg mice following immunization with NP-CGG were examined. As the result, there were modest reductions in the NP-specific IgM and IgG1 levels and the impairment was more prominent in the IgG2b and IgG2c levels in Ig $\beta$ -Tg mice after both primary and secondary immunizations (Fig. 4c). On day 21 post-immunization, the relative affinity of IgG1 for NP hapten was significantly reduced in Ig $\beta$ -Tg mice (Fig. 4d). Consistent with these data, the frequency of W33L replacement, which reflects the affinity maturation of the VH186.2 variable region to the NP hapten, among the over all mutations in the VH186.2 gene was reduced in Ig $\beta$ -Tg mice (Fig. 4e).

**Ig $\beta$  down-regulation in GC B cells is partly mediated by IL-21.** It has recently been reported that the absence of IL-21 signaling profoundly affects the T-dependent B cell response through the impairment of the GC reaction including antigen-specific IgG titer and affinity maturation<sup>24–26</sup>. To examine the possibility that IL-21 may exert its function by modulating the Ig $\beta$  level in GC B cells, it was first evaluated whether IL-21 has the potential to induce Ig $\beta$  down-regulation in purified B cells. Activation



**Figure 4.** Forced expression of Igβ in B cells impaired the antibody production and affinity maturation. **(a)** Expression of Igβ in B220<sup>+</sup>CD38<sup>-</sup>GL7<sup>-</sup> splenocytes from immunized Igβ-Tg (Igβ-Tg; line) and wild-type litter mates (wild type; shaded). **(b)** Expression of CD38 and GL7 in B220<sup>+</sup> splenocytes from immunized wild type litter mates (**top left**) and Igβ-Tg mice (**top right**). Cell cycle analysis of CD38<sup>-</sup>GL7<sup>-</sup> GC B cells from wild type litter mates (**bottom left**) and Igβ-Tg mice (**bottom right**). Numbers indicate the percentages of cells in each gate. **(c)** Serum anti-NP titers in each isotype after the immunization. Igβ-Tg (open circle) and wild type litter mates (closed circle) were immunized with NP-CGG on day 0 (1°) and boosted 42 d later (2°). Sera were collected at indicated day and assayed by ELISA as described in Methods section. Error bars indicate SEM for each group. Each group consists of 5 mice. Representative data from 3 repeated experiments is shown. **(d)** Affinity maturation of anti-NP IgG1 antibody in NP-CGG immunized Igβ-Tg (closed bar) and wild type litter mates (open bar). Error bars indicate SEM for each group. **(e)** Decreased frequency of somatic hypermutation in immunized Igβ-Tg mice. CD38<sup>-</sup>GL7<sup>+</sup>GC B cells were isolated from Igβ-Tg and wild type litter mates 21 days after the immunization. Total RNA was extracted and reverse transcribed to obtain cDNA samples. VH186.2 gene was then amplified and sequenced for the mutation analysis.



by IL-4 and an anti-CD40 antibody for 6 days, with or without IL-21, revealed that the frequency of Ig $\beta$  down-regulated cells increased in an IL-21 dose-dependent manner (Fig. 5a,b). When cell surface expression of CD138 was analyzed in these cells, up-regulation of CD138 was not detected in these Ig $\beta$  down-regulated B cells (Fig. S1c, right). These results indicate that IL-21 has the potential to directly act on activated B cells to regulate Ig $\beta$  expression.

To further examine the involvement of IL-21 in the down-regulation of Ig $\beta$  *in vivo*, the expression level of Ig $\beta$  in GC B cells from IL-21 receptor-deficient (IL-21R-KO) mice<sup>24</sup> was analyzed. As shown in Fig. 5c, the down-regulation of Ig $\beta$  was significantly impaired in the immunized IL-21R-KO mice. Furthermore, the frequency of CD38<sup>+</sup>GL7<sup>+</sup>B cells in the spleen from IL-21R-KO mice was reduced to approximately 50% as compared with that of wild type mice (0.51% to 0.25%) (Fig. 5d), accompanied by the reduction of proliferating (S + G2/M) Ig $\beta$ -Int cells (13.5% to 9.4%) (Fig. 5e).

Taken together along with the *in vitro* data, these results suggest that IL-21 may act on B cells directly or indirectly to down-regulate the expression of Ig $\beta$  in the GC, which may be required for the effective progression of the cell cycle in these cells and/or for differentiation to the proliferating population.

**Blocking of Ig $\beta$  down-regulation ameliorated autoimmune phenotype in BXS-B-Yaa mice.** It has been reported that IL-21 plays an essential role in the phenotypes of BXS-B-Yaa autoimmune-prone mice<sup>27,28</sup>. However, the precise mechanism of how IL-21 is involved in the generation and accumulation of autoreactive B cells has not been fully elucidated. To examine whether IL-21-mediated Ig $\beta$  down-regulation may contribute to the pathogenesis of BXS-B-Yaa mice, Ig $\beta$ -Tg mice were back-crossed to this strain. The percentage of Ig $\beta$ -Int cells among the B220<sup>+</sup>CD38<sup>+</sup>GL7<sup>+</sup> GC B cell population in BXS-B-Yaa mice was higher than in the wild type mice (56% to 41%), presumably due to the higher serum IL-21 level in BXS-B-Yaa mice (Fig. 6a)<sup>27,28</sup>. When Ig $\beta$ -Tg mice were back-crossed to BXS-B-Yaa, their percentage decreased from 56% to 18.6% and that of Ig $\beta$ -Hi cells increased reciprocally indicating that the Ig $\beta$  transgene is expressed as expected in the BXS-B-Yaa background (Fig. 6a). As shown in Fig. 6b,c, serum levels of anti-dsDNA IgG and proteinuria were ameliorated in BXS-B-Yaa mice carrying the Ig $\beta$  transgene. In addition, these mice tend to survive for a longer period compared with littermate BXS-B-Yaa mice (Fig. 6d). Assuming that autoantibody-producing cells are derived from GC B cells<sup>29</sup>, these results suggest that the modulation of Ig $\beta$  expression level affects not only positive selection, but also the accumulation of autoreactive B cells in the GC at least in the BXS-B-Yaa background.


## Discussion

In this study, we demonstrated quantitatively that the cell surface expression of Ig $\beta$  was more prominently lowered as compared with that of Ig $\alpha$  in the murine GC B cells and that the inhibition of Ig $\beta$  down-regulation resulted in the impairment of T-dependent immune responses. These results indicate that the down-regulation of Ig $\beta$  in GC B cells is essential for optimal antibody production and affinity maturation. Furthermore, crossing Ig $\beta$ -Tg mice with BXS-B-Yaa autoimmune-prone mice ameliorated the pathogenic phenotype.


Clonal selection of GC B cells with different antigen specificities and/or affinities that were raised by the somatic hypermutation resulted in affinity maturation<sup>11,30</sup>. During this selection stage, each B cell did not continue to retain BCRs with different specificities and/or affinities on its surface simultaneously, and could change their BCR specificity promptly to be selected<sup>31</sup>. This leads to the hypothesis that there may exist a machinery that excludes pre-existing BCRs with low affinity from the cell surface. Based on previous reports that the concomitant expression of Ig $\alpha$  and Ig $\beta$  is required for the surface expression of IgM<sup>4-6</sup>, it can be assumed that the down-regulation of Ig $\beta$  may contribute to the prompt clearance of surface IgM BCRs, most of which remain unmutated, in B cells that in turn start expressing mutated IgG in turn. Supporting this assumption, we have recently found that membrane-attached IgG1 can be expressed on the cell surface of GP2-293 cells and Ig $\beta$  down-regulated Ramos B cells, in contrast to IgM which could not be expressed on the surface of these cells<sup>32</sup>.

It has been reported that somatic hypermutation plays an essential roles in the generation of autoreactive antibody-producing B cells in some case of SLE<sup>33-35</sup>. However, it was also revealed in these studies that in addition to the pathogenic somatic hypermutation, failure in the selection stage was also required for the onset of the diseases. When Ig $\beta$ -Tg mice were crossed with BXS-B-Yaa mice, the pathogenesis observed in these mice was ameliorated. Although the precise molecular mechanism underlying this ameliorated phenotype remains to be elucidated, it may be possible that in the Ig $\beta$ -Tg background, class switching of GC B cells may proceed preferentially through the IgM<sup>+</sup>GL7<sup>+</sup>Ig $\beta$ -Hi population as in the case of *in vitro* class switching where expression of Ig $\beta$  is sustained at a comparable level as the naive B cells (Fig. S2). Under such condition, a skewed positive selection for the self-antigen or affinity maturation that may affect the autoimmune symptoms may be partly bypassed because down-regulation of Ig $\beta$  seems to be required for these processes to proceed efficiently.

Although the physiological relevance of Ig $\beta$  down-regulation is clear, it remains to be determined how the expression of Ig $\beta$  is modulated. In this study, it was revealed that Ig $\beta$  down-regulation in the GC B cells was modulated at least in part at the transcription level (Figs. 1c and 2d). Regarding the transcription control elements in the Ig $\beta$  promoter, three silencing motifs have been reported<sup>36,37</sup>. Among them, the most effective A+T-rich upstream sequence has been reported to bind Oct-1 and Oct-2<sup>37</sup>. Although these molecules are fascinating candidates for responsible transcription factors, the direct involvement



**Figure 5.** Impaired GC formation by inhibition of IL-21-mediated Igβ down-regulation. **(a)** Expression of Igβ in activated splenic B cells supplemented with IL-21. Splenocytes depleted of T cells were cultured with IL4 (10 ng/ml), and anti-CD40 antibody (1 μg/ml) supplemented with indicated concentrations of IL-21 for 6 days. Expression levels of Igβ and GL7 were analyzed by flow cytometric analysis. The numbers under the square gate indicate the percentages of Igβ down-regulated population in the gate. Representative data from 3 experiments is shown. **(b)** Average percentages and SEM of Igβ low cells from repeated experiments as in **(a)** were shown. **(c)** Flow cytometric analysis of Igβ expression in B220<sup>+</sup>CD38<sup>-</sup>GL7<sup>-</sup> splenocytes from immunized wild type (shaded) and IL-21R-deficient (line) mice. **(d)** Flow cytometric analysis of the splenic B cells (B220<sup>+</sup>) from wild type litter mates (**left**) and IL-21R-deficient mice (**right**) 10 days after the immunization with NP-CGG. The numbers indicate the percentage of CD38<sup>+</sup>GL7<sup>+</sup> cells within the gate. **(e)** Cell cycle analysis of B220<sup>+</sup>CD38<sup>-</sup>GL7<sup>-</sup> GC B cells from immunized IL-21R-deficient mice. Splenocytes from wild type litter mates (**left**) and IL21R-deficient mice (**right**) 10 days after the immunization were stained with Hoechst 33342 dye in combination with antibodies to Igβ.



**Figure 6.** Amelioration of auto-reactive antibody production and proteinuria by suppression of Igβ down-regulation in BXSB-Yaa autoimmune prone mice. Igβ-Tg mice were back-crossed to BXSB-Yaa for 9 generation and obtained male mice were subjected to further analyses. Serum titers in NZB and wild type C57BL/6 mice were also measured concomitantly as positive band negative controls, respectively. (a) Percentages of Igβ-Hi and Igβ-Int B cell population in B220<sup>+</sup>CD38<sup>+</sup>GL7<sup>+</sup> GC B cell subset from immunized C57BL/6 (open bars), BXSB-Yaa background wt (dark shaded bars) and Igβ-Tg (light shaded bars) mice are shown. (b) Serum titers of anti-dsDNA antibodies were measured in BXSB-Yaa background wt and Igβ-Tg mice at indicated ages. Each symbol indicate the serum levels for individual mouse. Thick lines indicate the average values and thin lines indicate SEM values. (c) Proteinuria was measured at 20 weeks of age. Each circle indicate the serum levels for individual mouse. (d) Survival curve of BXSB-Yaa background wt and Igβ-Tg mice.

of Oct-1 and/or Oct-2 in the regulatory mechanisms of Ig $\beta$  down-regulation in GC B cells remains to be tested.

With regard to soluble factors that may modulate Ig $\beta$  expression, it was also revealed that Ig $\beta$  down-regulation is partly mediated by IL-21 (Fig. 5). In accordance with these results, IL-21R-deficient mice showed impairment in the development of GC B cells and in affinity maturation<sup>25,26,38–40</sup>. However, because the blockade of Ig $\beta$  down-regulation was not complete in IL-21R-deficient mice, it is evident that IL-21-independent machinery also exists. Antigen receptors of GC B cells are believed to be cross-linked by the cognate antigen during positive selection, which may result in the internalization of the BCR with its associating molecules, including Ig $\alpha$  and Ig $\beta$ . Thus, in addition to transcriptional regulation, we cannot completely rule out the possibility that post-translational modulation such as internalization, circulation or degradation of Ig $\beta$  in association with BCRs may also be involved in its down-regulation<sup>15–17,41,42</sup>. Indeed, existence of an as-yet-unknown mechanism for post-transcriptional modulation of Ig $\beta$  expression, which differ from Ig $\alpha$ , was suggested in human GC B cells<sup>17</sup>. The surface expression levels of Ig $\alpha$  were not dramatically down-regulated in the B220<sup>+</sup>CD38<sup>+</sup>GL7<sup>+</sup> GC B cell population as compared with that of the B220<sup>+</sup>CD38<sup>+</sup>GL7<sup>+</sup> B cells, in contrast to the expression levels of Ig $\beta$ . However, a population that was up-regulating the surface Ig $\alpha$  level could be observed, suggesting that still-unknown, unique mechanism exists for the regulation of Ig $\alpha$ . Despite the marginal down-regulation of Ig $\alpha$  in GC B cells, a decrease in the Ig $\alpha$  mRNA level in GC B cells was detected (data not shown). This also suggests the existence of post-transcriptional regulation in the case of Ig $\alpha$ . However, as aforementioned, a simple explanation of our results mainly by endocytosis of the IgM BCR complex associating with Ig $\alpha$ /Ig $\beta$  heterodimer seems unlikely.

## Methods

**Mouse and immunization.** Ig $\beta$ -Tg mouse was generated using pEMV4C8-Ig $\beta$ , which consists of mouse E $\mu$  enhancer, mouse VH4C8 promoter, rabbit  $\beta$ -globin intron followed by HA-tagged mouse Ig $\beta$  cDNA and rabbit  $\beta$ -globin derived polyA signal. Expression vector was a generous gift from Dr. Takeshi Watanabe. Linearized expression vector was microinjected to C57BL/6 blastocyst and 2 founder lines, which both possess 3 copies of transgene, were established (Fig. S3) (PhoenixBio). IL-21 receptor-deficient mice<sup>24</sup> was provided from Drs. Warren J. Leonard and Katsutoshi Ozaki. Immunizations were carried out as described previously<sup>43</sup>. In brief, 8–12 weeks old C57BL/6 mice were injected i.p. with 100ug NP-CGG (Biosearch Technologies) adsorbed to 3 mg Imject Alum (Thermo Fisher Scientific) according to the manufacturer's instructions. All experimental protocols were approved by animal research committee in the Graduate School of Medicine, Kyoto University and carried out in accordance with the guidelines for the animal experiments in Kyoto University.

**Flow cytometry and histological analysis.** All cells were stained with indicated combinations of antibodies, and analyzed with FACSaria II (BD Biosciences). Cells were stained in combination with propidium iodide (Sigma) to exclude dead cells as described previously<sup>43</sup>.

Frozen 8  $\mu$ m sections of spleens from immunized mice were subjected to immunohistochemical staining. In brief, sections were fixed with 1:1 mixture of methanol (Wako) and acetone (Wako), blocked with 1% BSA in PBS for 1 h, and stained with indicated antibodies in PBS with 0.1% BSA for 1 h. Finally, the sections were washed and mounted with SlowFade Gold (Invitrogen). Stained sections were visualized using LSM 710 NLO confocal fluorescent microscope with ZEN software (ZEISS).

**Quantitative RT-PCR and PCR for cloning.** Total RNA from sorted splenic cells was extracted using TRIzol Reagent (Invitrogen). Then, cDNA was synthesized from total RNA using SuperScript III First-Strand Synthesis System (Invitrogen). Transcription levels of Ig $\beta$ , Bcl-6, AID, germline transcript from I $\mu$  locus (GLT- $\mu$ ), germline transcript from I $\gamma$ 1 locus (GLT- $\gamma$ 1) and postswitch transcript of C $\gamma$ 1 were determined by quantitative RT-PCR using the LightCycler 480 SYBR Green I Master and LightCycler 480 system (Roche). Relative expression levels of mRNA were calculated compared to those of  $\beta$ -actin.

**ELISA.** Serum anti-NP antibody levels were measured by ELISA as described below. 96-well flat-bottom plates (Thermo Fisher Scientific) were coated with 10  $\mu$ g/ml NP<sub>54</sub>-BSA (Biosearch Technologies), and then the wells were blocked with 5-fold diluted Blocking-One (Nacalai Tesque). Appropriately diluted serum samples were added to the wells and incubated at room temperature for 2 h. The plates were washed and incubated for another 1 h with diluted anti-IgM, -IgG1, -IgG2c and -IgG2b antibodies conjugated with horse radish peroxidase (Bethyl Laboratories), respectively. Bound secondary antibody was assessed by incubation with ABTS substrate (Sigma-Aldrich). Absorbance at 405 nm (A405) in each well was measured.

For the analysis of affinity maturation, appropriately diluted serum samples were assayed using an NP<sub>4</sub>-BSA (Biosearch Technologies) -coated 96-well plate and an NP<sub>54</sub>-BSA-coated plate. The A405(NP<sub>4</sub>)/A405(NP<sub>54</sub>) ratio was calculated as relative affinity.

**Generation of polyclonal antibody to cell surface Ig $\alpha$ .** Synthetic peptide corresponding to 89 to 99 amino acids with a extra cystein residue at N-terminal (CPEVNKNHTRGLY) was conjugated to KLH

Antibody	Manufacturer	clone
anti-B220-APC-Cy7	BD Biosciences	RA3-6B2
anti-CD38-PE	eBioscience	90
anti-GL7-Alexa Fluor 647	eBioscience	GL-7
anti-CD79b-FITC	BD Biosciences	HM79b
anti-Fas-PE-Cy7	eBioscience	Jo2
anti-IgM-PE-Cy7	eBioscience	II/41
streptavidin-PE-Cy7	eBioscience	-
streptavidin-Texas Red	BD Biosciences	-
anti-IgG1	BD Biosciences	A85-1
anti-IgG1-Pacific Blue	BD Biosciences	A85-1
anti-CD79b-biotin	BD Biosciences	HM79b
PNA-FITC	Seikagaku Kogyo	-

**Table 1.** List of antibodies used in the experiments

Name of the oligo	Sequence of the oligo
Ig $\beta$ -qPCRf	CGAGGTTTGACGCCAAAAAG
Ig $\beta$ -qPCRr	CACAATGCGTCCCTCTTCTG
Bcl-6-qPCRf	ATCTGTGGCACTCGCTTCC
Bcl-6-qPCRr	AGTCGCAGTTGGCTTTTGTG
AID-qPCRf	AAGCCTGGGAAGGGCTACA
AID-qPCRr	GACATTCCAGGAGGTTGCTTTC
GLT $\mu$ -qPCRf	CTCTGGCCCTGCTTATTGTTG
GLT $\mu$ -qPCRr	GAAGACATTGGGAAGGACTGACT
GLT $\gamma$ 1-qPCRf	AGGAATGTGTTTGGCATGGAC
GLT $\gamma$ 1-qPCRr	CACTGTCACTGGCTCAGGGAA
postswitch C $\gamma$ 1-qPCRf	CTCTGGCCCTGCTTATTGTTG
postswitch C $\gamma$ 1-qPCRr	CACTGTCACTGGCTCAGGGAA
$\beta$ -actin-qPCRf	GGCTGTATTCCCCTCCATCG
$\beta$ -actin-qPCRr	CCAGTTGGTAACAATGCCATGT
VH186.2F	TTCTTGGCAGCAACAGCTACA
IgG1R	CACTGTCACTGGCTCAGGGAA

**Table 2.** List of the oligos used in the experiments

and repeatedly immunized to the rabbits and the obtained anti-serum was affinity purified with the immunized peptide conjugated to BSA (Scrum). Specificity of the antibody was confirmed by western blot analysis of murine splenic B cells and surface staining of NIH/3T3 cells transfected with Ig $\alpha$ , Ig $\beta$  and single chain EGFP-Fc $\gamma$ 1. Also it was confirmed that no other spleen cells except B220<sup>+</sup> cells were stained with the antibody.

**Analysis of somatic hypermutation.** Total RNA from the sorted splenocytes of immunized C57BL/6 or Ig $\beta$ -Tg mice was extracted and reverse transcribed to obtain cDNA, which was subjected to PCR using KOD DNA polymerase (Toyobo) with VH186.2-specific VH186.2F and IgG1-specific IgG1R primers and cloned into the plasmid PCR-blunt TOPO (Invitrogen) to analyze the sequence of the VH gene.

**Measurement of autoantibody and proteinuria.** Serum anti-dsDNA titers were measured using mouse anti-dsDNA IgG ELISA kit (Alpha Diagnostic International Inc.) according to the manufacturer's instruction. Proteinuria was measured using Albustix Reagent Strips (SIEMENS) according to the manufacturer's instruction.



**Antibodies and oligo nucleotides.** Antibodies and oligo nucleotides used in this study were listed in Table 1 and Table 2, respectively.

## References

- Reth, M. Antigen receptors on B lymphocytes. *Annu. Rev. Immunol.* **10**, 97–121, DOI:10.1146/annurev.iy.10.040192.000525 (1992).
- Reth, M. & Wienands, J. Initiation and processing of signals from the B cell antigen receptor. *Annu. Rev. Immunol.* **15**, 453–479, DOI:10.1146/annurev.immunol.15.1.453 (1997).
- Kurosaki, T., Shinohara, H. & Baba, Y. B cell signaling and fate decision. *Annu. Rev. Immunol.* **28**, 21–55, DOI:10.1146/annurev.immunol.021908.132541 (2010).
- Hombach, J., Tsubata, T., Leclercq, L., Stappert, H. & Reth, M. Molecular components of the B-cell antigen receptor complex of the IgM class. *Nature* **343**, 760–762, DOI:10.1038/343760a0 (1990).
- Matsuuchi, L. *et al.* The membrane IgM-associated proteins MB-1 and Ig-b are sufficient to promote surface expression of a partially functional B-cell antigen receptor in a nonlymphoid cell line. *Proc. Natl. Acad. Sci. USA* **89**, 3404–3408 (1992).
- Kitamura, D., Roes, J., Kuhn, R. & Rajewsky, K. A B cell-deficient mouse by targeted disruption of the membrane exon of the immunoglobulin m chain gene. *Nature* **350**, 423–426, DOI:10.1038/350423a0 (1991).
- Lam, K. P., Kuhn, R. & Rajewsky, K. *In vivo* ablation of surface immunoglobulin on mature B cells by inducible gene targeting results in rapid cell death. *Cell* **90**, 1073–1083, DOI:S0092-8674(00)80373-6 (1997).
- Gong, S. & Nussenzweig, M. C. Regulation of an early developmental checkpoint in the B cell pathway by Igb. *Science* **272**, 411–414 (1996).
- Kraus, M. *et al.* Interference with immunoglobulin (Ig) a immunoreceptor tyrosine-based activation motif (ITAM) phosphorylation modulates or blocks B cell development, depending on the availability of an Igb cytoplasmic tail. *J. Exp. Med.* **194**, 455–469 (2001).
- Kraus, M., Alimzhanov, M. B., Rajewsky, N. & Rajewsky, K. Survival of resting mature B lymphocytes depends on BCR signaling via the Igα/β heterodimer. *Cell* **117**, 787–800, DOI:10.1016/j.cell.2004.05.014 S0092867404005306 (2004).
- Rajewsky, K. Clonal selection and learning in the antibody system. *Nature* **381**, 751–758, DOI:10.1038/381751a0 (1996).
- Nutt, S. L. & Tarlinton, D. M. Germinal center B and follicular helper T cells: siblings, cousins or just good friends? *Nat. Immunol.* **12**, 472–477 (2011).
- MacLennan, I. C. Germinal centers. *Annu. Rev. Immunol.* **12**, 117–139, DOI:10.1146/annurev.iy.12.040194.001001 (1994).
- Allen, C. D., Okada, T. & Cyster, J. G. Germinal-center organization and cellular dynamics. *Immunity* **27**, 190–202, DOI:10.1016/j.immuni.2007.07.009 (2007).
- Khalil, A. M., Cambier, J. C. & Shlomchik, M. J. B cell receptor signal transduction in the GC is short-circuited by high phosphatase activity. *Science* **336**, 1178–1181, DOI:10.1126/science.1213368 (2012).
- Guo, B. & Rothstein, T. L. IL-4 upregulates Iga and Igb protein, resulting in augmented IgM maturation and B cell receptor-triggered B cell activation. *J. Immunol.* **191**, 670–677, DOI:10.4049/jimmunol.1203211 (2013).
- Huang, X. *et al.* Downregulation of the B-cell receptor signaling component CD79b in plasma cell myeloma: a possible post transcriptional regulation. *Pathol. Int.* **61**, 122–129, DOI:10.1111/j.1440-1827.2010.02634.x (2011).
- Smith, K. G., Nossal, G. J. & Tarlinton, D. M. FAS is highly expressed in the germinal center but is not required for regulation of the B-cell response to antigen. *Proc. Natl. Acad. Sci. USA* **92**, 11628–11632 (1995).
- Hao, Z. *et al.* Fas receptor expression in germinal-center B cells is essential for T and B lymphocyte homeostasis. *Immunity* **29**, 615–627, DOI:S1074-7613(08)00409-3 10.1016/j.immuni.2008.07.016 (2008).
- Good-Jacobson, K. L. *et al.* PD-1 regulates germinal center B cell survival and the formation and affinity of long-lived plasma cells. *Nat. Immunol.* **11**, 535–542, DOI:ni.1877 10.1038/ni.1877 (2010).
- Li, S. C. *et al.* Expression of Im-Cg hybrid germline transcripts subsequent to immunoglobulin heavy chain class switching. *Int. Immunol.* **6**, 491–497 (1994).
- Stavnezer, J., Guikema, J. E. & Schrader, C. E. Mechanism and regulation of class switch recombination. *Annu. Rev. Immunol.* **26**, 261–292, DOI:10.1146/annurev.immunol.26.021607.090248 (2008).
- Nojima, T. *et al.* *In vitro* derived germinal centre B cells differentially generate memory B or plasma cells *in vivo*. *Nat. Commun.* **2**, 465, DOI:ncomms1475 10.1038/ncomms1475 (2011).
- Ozaki, K. *et al.* A critical role for IL-21 in regulating immunoglobulin production. *Science* **298**, 1630–1634, DOI:10.1126/science.1077002 298/5598/1630 (2002).
- Linterman, M. A. *et al.* IL-21 acts directly on B cells to regulate Bcl-6 expression and germinal center responses. *J. Exp. Med.* **207**, 353–363, DOI:jem.20091738 10.1084/jem.20091738 (2010).
- Zotos, D. *et al.* IL-21 regulates germinal center B cell differentiation and proliferation through a B cell-intrinsic mechanism. *J. Exp. Med.* **207**, 365–378, DOI:jem.20091777 10.1084/jem.20091777 (2010).
- Ozaki, K. *et al.* Regulation of B cell differentiation and plasma cell generation by IL-21, a novel inducer of Blimp-1 and Bcl-6. *J. Immunol.* **173**, 5361–5371 (2004).
- Bubier, J. A. *et al.* A critical role for IL-21 receptor signaling in the pathogenesis of systemic lupus erythematosus in BXSB-Yaa mice. *Proc. Natl. Acad. Sci. USA* **106**, 1518–1523, DOI:0807309106 10.1073/pnas.0807309106 (2009).
- Moisini, I. *et al.* The Yaa locus and IFN-α fine-tune germinal center B cell selection in murine systemic lupus erythematosus. *J. Immunol.* **189**, 4305–4312, DOI:10.4049/jimmunol.1200745 (2012).
- Goodnow, C. C., Vinuesa, C. G., Randall, K. L., Mackay, F. & Brink, R. Control systems and decision making for antibody production. *Nat. Immunol.* **11**, 681–688, DOI:ni.1900 10.1038/ni.1900 (2010).
- Tarlinton, D. M. & Smith, K. G. Dissecting affinity maturation: a model explaining selection of antibody-forming cells and memory B cells in the germinal centre. *Immunol. Today* **21**, 436–441, DOI:S016756990001687X (2000).
- Todo, K., Koga, O., Nishikawa, M. & Hikida, M. IgG1 cytoplasmic tail is essential for cell surface expression in Igβ down-regulated cells. *Biochem. Biophys. Res. Commun.* **445**, 572–577, DOI:10.1016/j.bbrc.2014.02.037 (2014).
- Detanico, T. *et al.* Somatic mutagenesis in autoimmunity. *Autoimmunity* **46**, 102–114, DOI:10.3109/08916934.2012.757597 (2013).
- Guo, W. *et al.* Somatic hypermutation as a generator of antinuclear antibodies in a murine model of systemic autoimmunity. *J. Exp. Med.* **207**, 2225–2237, DOI:10.1084/jem.20092712 (2010).
- Zhang, J., Jacobi, A. M., Wang, T. & Diamond, B. Pathogenic autoantibodies in systemic lupus erythematosus are derived from both self-reactive and non-self-reactive B cells. *Molecular medicine* **14**, 675–681, DOI:10.2119/2008-00066.Zhang (2008).
- Malone, C. S., Omori, S. A. & Wall, R. Silencer elements controlling the B29 (Igb) promoter are neither promoter- nor cell-type-specific. *Proc. Natl. Acad. Sci. USA* **94**, 12314–12319 (1997).
- Malone, C. S., Patrone, L., Buchanan, K. L., Webb, C. F. & Wall, R. An upstream Oct-1- and Oct-2-binding silencer governs B29 (Igb) gene expression. *J. Immunol.* **164**, 2550–2556, DOI:ji\_v164n5p2550 (2000).

38. Fazilleau, N., Mark, L., McHeyzer-Williams, L. J. & McHeyzer-Williams, M. G. Follicular helper T cells: lineage and location. *Immunity* **30**, 324–335, DOI:S1074-7613(09)00114-9 10.1016/j.immuni.2009.03.003 (2009).
39. King, C., Tangye, S. G. & Mackay, C. R. T follicular helper (TFH) cells in normal and dysregulated immune responses. *Annu. Rev. Immunol.* **26**, 741–766, DOI:10.1146/annurev.immunol.26.021607.090344 (2008).
40. Vinuesa, C. G., Tangye, S. G., Moser, B. & Mackay, C. R. Follicular B helper T cells in antibody responses and autoimmunity. *Nat. Rev. Immunol.* **5**, 853–865, DOI:nri1714 10.1038/nri1714 (2005).
41. Yuseff, M. I., Lankar, D. & Lennon-Dumenil, A. M. Dynamics of membrane trafficking downstream of B and T cell receptor engagement: impact on immune synapses. *Traffic* **10**, 629–636, DOI:TRA913 10.1111/j.1600-0854.2009.00913.x (2009).
42. Zhang, M. *et al.* Ubiquitinylation of Ig $\beta$  dictates the endocytic fate of the B cell antigen receptor. *J. Immunol.* **179**, 4435–4443, DOI:179/7/4435 (2007).
43. Hikida, M. *et al.* PLC- $\gamma$ 2 is essential for formation and maintenance of memory B cells. *J. Exp. Med.* **206**, 681–689, DOI:10.1084/jem.20082100 (2009).

## Acknowledgement

We thank Drs. Warren J. Leonard and Katsutoshi Ozaki for generously providing IL-21 receptor-deficient mice, Dr. Takeshi Watanabe for critical reading of the manuscript.

## Author Contributions

K.T. designed research, did experiments, analyzed data and wrote the manuscript; M.H. designed research, did experiments, analyzed data and wrote the manuscript; O.K. and M.N. did the experiments.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>


**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Todo, K. *et al.* Modulation of Ig $\beta$  is essential for the B cell selection in germinal center. *Sci. Rep.* **5**, 10303; doi: 10.1038/srep10303 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

# SCIENTIFIC REPORTS



## **Erratum:** Modulation of Ig $\beta$ is essential for the B cell selection in germinal center

Kagefumi Todo, Orié Koga, Miwako Nishikawa & Masaki Hikida

*Scientific Reports* 5:10303; doi: 10.1038/srep10303; published online 18 May 2015; updated on 05 August 2015

The Competing financial interests statement in this Article should read:

“This work was supported in part by Astellas Pharma Inc. in the Formation of Innovation Center for Fusion of Advanced Technologies Program.”